



## Influence of Nutrients involved in One-Carbon Metabolism on DNA Methylation in Adults

Amenyah, S., Hughes, C., Ward, M., Roxborough, S., Deane, J., Thursby, S-J., Walsh, CP., Kok, D., Strain, S., McNulty, H., & Lees Murdock, D. (2020). Influence of Nutrients involved in One-Carbon Metabolism on DNA Methylation in Adults: A Systematic Review and Meta-Analysis. *Nutrition Reviews*, 78(8), 647-666. [nuz094]. <https://doi.org/10.1093/nutrit/nuz094>

[Link to publication record in Ulster University Research Portal](#)

**Published in:**  
Nutrition Reviews

**Publication Status:**  
Published (in print/issue): 31/08/2020

**DOI:**  
[10.1093/nutrit/nuz094](https://doi.org/10.1093/nutrit/nuz094)

**Document Version**  
Author Accepted version

### General rights

Copyright for the publications made accessible via Ulster University's Research Portal is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

### Take down policy

The Research Portal is Ulster University's institutional repository that provides access to Ulster's research outputs. Every effort has been made to ensure that content in the Research Portal does not infringe any person's rights, or applicable UK laws. If you discover content in the Research Portal that you believe breaches copyright or violates any law, please contact [pure-support@ulster.ac.uk](mailto:pure-support@ulster.ac.uk).

**Article type: Lead article** (Systematic review and meta-analysis)

**Title: Influence of Nutrients involved in One-Carbon Metabolism on DNA Methylation in Adults - A Systematic Review and Meta-Analysis**

**Author names:** Sophia D. Amenyah<sup>1,2</sup>, Catherine F. Hughes<sup>2</sup>, Mary Ward<sup>2</sup>, Samuel Rosborough<sup>1</sup>, Jennifer Deane<sup>1</sup>, Sara-Jayne Thursby<sup>1</sup>, Colum P. Walsh<sup>1</sup>, Dieuwertje E. Kok<sup>3</sup>, JJ Strain<sup>2</sup>, Helene McNulty<sup>2</sup>, Diane J. Lees-Murdock<sup>1</sup>

**Author Affiliations:** <sup>1</sup>Genomic Medicine Research Group (SDA, SR, JD, SJ-T, CPW and DLM) and <sup>2</sup>Nutrition Innovation Centre for Food and Health (NICHE; SDA, SFH, MW, JJS and HM), School of Biomedical Sciences, Ulster University, Coleraine, Northern Ireland, United Kingdom and <sup>3</sup>Division of Human Nutrition and Health, Wageningen University, PO Box 17, 6700 AA, Wageningen, The Netherlands (DEK).

**Authors' last names:** Amenyah, Hughes, Rosborough, Deane, Thursby, Walsh, Kok, Strain, McNulty, Ward, Lees-Murdock

**Corresponding Author:** Dr Diane Lees-Murdock, School of Biomedical Sciences, Ulster University, Cromore Road, Coleraine, Northern Ireland, UK. BT52 1SA. Tel: +44 (0) 28 7012 3166, E-mail: dj.lees@ulster.ac.uk

**Abbreviations used:** CI, confidence interval; CVD, cardiovascular disease; DMP, differentially methylated position; DMR, differentially methylated region; LC-MS, liquid chromatography-tandem mass spectrometry; LINE-1, long interspersed nuclear elements; MTHFR, methylenetetrahydrofolate reductase; RBC, red blood cell; RCT, randomized controlled trial; SAM, S-adenosylmethionine; SD, standard deviation; SEM, standard error of mean; TSS, transcription start site; UTR, untranslated region.

## ABSTRACT

**Context:** Aberrant DNA methylation is linked to various diseases. The supply of methyl groups for methylation reactions is mediated via S-adenosylmethionine which depends on the availability of folate and related B-vitamins.

**Objectives:** To investigate the influence of key nutrients involved in one-carbon metabolism on DNA methylation in adults.

**Data sources:** Systematic literature searches were conducted in the Cochrane library, Medline, Embase, CINAHL Plus, Scopus and Web of Science databases. Studies that met the inclusion criteria and were published in English were included.

**Data extraction:** The first author, study design, sample size, population characteristics, type of intervention and duration, tissue type or cells analyzed, molecular techniques and DNA methylation outcomes.

**Data synthesis:** A meta-analysis of RCTs was conducted to investigate the effect of one-carbon metabolism nutrients on global DNA methylation. Functional analysis and visualization was performed using BioVenn software.

**Results:** From a total of 2620 papers screened by title, 53 studies met the inclusion criteria. Qualitative analysis indicates significant associations between one-carbon metabolism nutrients and DNA methylation. In meta-analysis of RCTs stratified by method of laboratory analysis, supplementation with folic acid alone or in combination with vitamin B-12 significantly increased global DNA methylation in studies employing LC-MS, which had markedly lower heterogeneity ( $n = 3$ ,  $Z = 3.31$ ,  $P = 0.0009$ ;  $I^2 = 0\%$ ) in comparison to other methods. Functional analysis highlighted a subset of 12 differentially methylated regions that were significantly related to both folate and vitamin B-12 biomarkers.

46 **Conclusions:** This study supports significant associations between one-carbon metabolism  
47 nutrients and DNA methylation. However, standardization of DNA methylation techniques is  
48 recommended to reduce heterogeneity and facilitate comparison across studies.

49 **Systematic Review registration:** PROSPERO registration number: CRD42018091898.

50 **Key words:** One-carbon metabolism nutrients, DNA methylation, B-vitamins, one-carbon  
51 metabolism, systematic review, meta-analysis

## INTRODUCTION

DNA methylation is the most stable epigenetic mechanism in mammals. It is important in the regulation of gene expression and maintaining genome stability both locally and at the global level <sup>1,2</sup>. Changes in methylation occur in utero or in early life and are subject to age-related changes during an organism's lifetime <sup>3-6</sup>. This systematic review focuses on methylation changes during adult life. Aberrant DNA methylation in adults has been linked to aging and implicated in many diseases including cancer and cardiovascular disease <sup>1,2,4</sup>. Understanding the role of B-vitamins in regulating DNA methylation in aging and their roles in disease pathophysiology is essential in both the diagnosis and treatment of many diseases <sup>7</sup>.

DNA methylation has been shown to be responsive to environmental shifts such as changes in diet or nutritional status <sup>8,9</sup>. Through the interaction with nutrients involved in one-carbon metabolism, methylation of specific genes can be modified, influencing gene expression and phenotypes <sup>10-12</sup>. Additionally, nutritional status can interact with specific genetic variants of key genes in one-carbon metabolism to modulate health offering a unique opportunity for dietary based interventions that target diseases linked to altered DNA methylation <sup>13,14</sup>.

One-carbon metabolism is one of the main metabolic networks by which nutrients interact biologically to modulate DNA methylation. Nutrients involved in one-carbon metabolism include folate, vitamin B-12, vitamin B-6, riboflavin (vitamin B-2), choline, betaine, methionine and homocysteine <sup>15</sup>. Folate and related B-vitamins provide the substrates and cofactors to ensure the efficient functioning of one-carbon metabolism <sup>16-18</sup>. Of particular note, the folate and methionine pathways in one-carbon metabolism generate S-adenosylmethionine (SAM), the universal methyl donor, required for numerous biological reactions including DNA, RNA and histone methylation.

Currently, evidence for the role of specific nutrients within the network on DNA methylation is conflicting. In several studies, intervention with B-vitamins, mainly folic acid, led to alterations in global, gene-specific or CpG site-specific DNA methylation<sup>19–21</sup>, however conversely, other studies report no changes in methylation in response to folic acid or B-vitamin supplementation<sup>22–24</sup>. Furthermore, very little is known about doses, dietary exposure levels or extent of depletion necessary to elicit these epigenetic changes. Additionally, conditions such as life stage or health status at which one-carbon metabolism related nutrients have the largest modulatory effects on DNA methylation are currently not fully understood. There is therefore a need to systematically analyze and evaluate the current evidence for the influence of relevant nutrients involved in one-carbon metabolism on DNA methylation.

The aim of this study was to conduct a systematic review to investigate the influence of nutrients involved in one-carbon metabolism on DNA methylation in adult populations. In addition, a meta-analysis of RCTs was conducted to examine the effects of supplementation with relevant nutrients on global DNA methylation.

## **METHODS**

This systematic review was conducted according to PRISMA guidelines (PRISMA checklist provided in **Supplementary Table S1**) and a registered protocol (PROSPERO 2018, CRD42018091898). Screening of eligible studies, full-text assessment, data extraction and quality assessment of studies was independently carried out by two authors, discrepancies were discussed and resolved by consensus and where necessary moderated by a third reviewer. Studies were selected in accordance with the PICOS (population, intervention, comparison, outcome, and study design) criteria shown in **Table 1**.

### **Search Strategy and Study Selection**

Systematic literature searches were conducted in the Cochrane library, Medline (Ovid), Embase, CINAHL Plus, Scopus and Web of Science databases without any language restrictions in March 2019 (detailed search strategy provided in **Supplementary Table S2**). The full search strategy for all the searches combined terms related to one-carbon metabolism nutrients or synonyms (e.g. folate, vitamin B-12, riboflavin and vitamin B-6), DNA methylation (e.g. global, gene-specific, genome-wide methylation) and homocysteine are presented in **Supplementary Table S2**. Medical subject headings and key word searches were conducted in Embase, Medline, CINAHL Plus and Cochrane databases while searches in Scopus and Web of Science were carried out using only key word searches.

Following removal of duplicates, the titles and abstracts of studies retrieved from the literature search were screened for potentially eligible studies. Full text articles of potentially relevant articles were further reviewed using a pre-designed in/out form which included questions to assess each study's relevance for the review. Studies were considered eligible if they were original peer-reviewed full-text articles published in English and included all the defined outcomes.

## **Inclusion and Exclusion Criteria**

Studies conducted in adult humans investigating all of the following: 1) DNA methylation (global, gene-specific and genome-wide methylation), 2) nutrients involved in one-carbon metabolism and 3) circulating homocysteine levels (potential biomarker of one-carbon metabolism) were included in the current review. Studies involving 1) pregnant women and children, 2) *in-vitro* studies using human or animal cell lines and 3) studies conducted in animals were excluded from the analysis.

## **Data Extraction, Synthesis and Analysis**

Data extraction was carried out using a predesigned data collection sheet to extract relevant information from the selected studies. Information extracted included the name of

first author, study design, sample size, population characteristics, type of intervention and duration (intervention studies and randomized controlled trials), type of tissues or cells analyzed, molecular techniques and outcomes related to DNA methylation.

A narrative synthesis using descriptive statistics such as frequencies and percentages are presented for all studies included. The effects of supplementation with nutrients involved in one-carbon metabolism on DNA methylation are reported for RCTs and intervention studies. A meta-analysis examining the effect of supplementation with nutrients involved in one-carbon metabolism on global DNA methylation is included for RCT studies. Associations between one-carbon metabolism nutrients and DNA methylation are reported for observational studies. Owing to the considerable heterogeneity in study aims, designs and evaluated outcomes in the observational and intervention studies included, no quantitative analysis could be carried out for these type of studies.

### **Assessment of Risk of Bias**

Risk of bias of RCTs and intervention studies was assessed using the following key criteria: random sequence generation, allocation concealment, blinding of participants and outcome, incomplete outcome data, selective reporting and other sources of bias in accordance to the Cochrane Risk of Bias Assessment tool <sup>25</sup>. The risk of bias in each study was classified as low risk, high risk or unclear risk (either a lack of information or uncertainty over potential bias). Risk of bias of observational studies were assessed for key criteria: selection, comparability and outcome using the Newcastle-Ottawa scale<sup>26</sup>.

### **Quality of Reporting Studies**

Quality of reporting the studies included in the review was assessed using the STROBE (STrengthening the Reporting of OBservational studies in Epidemiology) checklist <sup>27</sup> for observational studies, the CONSORT (Consolidated Standards of Reporting Trials) checklist <sup>28</sup> for RCTs and a modification of the TREND statement <sup>29</sup> for intervention studies



without randomization. All questions on the appropriate checklists were considered for the studies included. The final score for each study was based on adherence to appropriate checklist criteria. A percentage score was calculated as the number of checklist criteria adhered to divided by the total number of questions on the checklist.

## **Meta-Analysis of the Effects of Supplementation with Nutrients involved One-Carbon Metabolism on Global DNA Methylation**

Meta-analysis of RCTs included in the review was conducted to examine the effects of supplementation with nutrients involved in one-carbon metabolism on global DNA methylation. No quantitative analysis could be carried out for the RCTs focusing on gene-specific methylation owing to the diverse range of candidate loci examined and resulting paucity of data for each target. Similarly, only 1 genome-wide methylation study was returned from the search preventing its inclusion in the meta-analysis. The remaining RCTs investigating global methylation were considered for inclusion in the meta-analysis only if they included a placebo or control group.

Data synthesis for meta-analysis was conducted using the standardized EURECA guidelines<sup>30</sup>. In cases where 2 publications reported data from the same study, they were linked and treated as one “main” intervention study. With the use of this approach, Pufulete *et al*<sup>19</sup> and Al-Ghnaniem Abbadi *et al.*<sup>22</sup> were treated as one study and the global methylation results reported in Al-Ghnaniem Abbadi were excluded from the meta-analysis. Further LINE-1 methylation data from Obeid *et al.*<sup>20</sup> and Pusceddu *et al.*<sup>31</sup> were from the same study and hence the data reported in Obeid *et al.*<sup>20</sup> were used in the meta-analysis. Where studies used more than one intervention strategy with one common placebo group<sup>32</sup>, each intervention arm and placebo group were treated as an independent study in the meta-analysis. Further, in studies where DNA methylation was measured at two different time

points post-supplementation<sup>23</sup> or in different tissues<sup>19</sup> in the same study, the results were treated as independent studies in the meta-analysis.

## Statistical Analysis

Review Manager 5.3 software (Cochrane Collaboration, 2014) was used to perform the meta-analysis. Mean methylation values and corresponding standard deviations were extracted from the included studies. In studies where the measure of variance was reported as SEM or CI<sup>19,23,24,33,34</sup>, the SD was estimated using Cochrane formulas<sup>35</sup>. The overall pooled effect (Z) was analyzed using the standardized mean difference and the random effects model. The random effects model estimates the between-study variance and uses this estimate to modify the weights assigned to individual studies when calculating the overall effect<sup>36</sup>. In order to establish if methylation is confounded by heterogeneity in one-carbon metabolism nutrient supplemented, DNA methylation technique or tissue analyzed, pre-specified subgroup and sensitivity analyses was carried out for each of those variables. Data are expressed as standardized mean difference (95% CI) and the overall effect Z (*P*-value).

Statistical heterogeneity was evaluated using chi square value, heterogeneity index ( $I^2$ ) statistics and corresponding *P*-value. Heterogeneity thresholds were defined according to Cochrane guidelines, with  $I^2$  between 0 - 40% indicative of low heterogeneity,  $I^2$  between 30 - 60% representing moderate heterogeneity,  $I^2$  between 50 - 90% representing substantial heterogeneity and  $I^2$  between 75 - 100% indicating considerable heterogeneity<sup>35</sup>. Potential publication bias for each study included in the meta-analysis was assessed by visual inspection of funnel plots and Egger's regression test<sup>37</sup>.

## Functional Analysis of Epigenome-wide Methylation

Functional analysis was carried out using differentially methylated regions (DMRs) previously identified to be related to both serum folate and vitamin B-12 in epigenome-wide methylation studies<sup>21</sup>. The DMRs were identified using the DMRcate package available

through Bioconductor in R statistical environment. Overlapping DMRs associated with serum folate or vitamin B-12 levels were visualized using BioVenn Software <sup>38</sup>.

## RESULTS

A total of 2620 records were identified through searches in 6 databases. After screening and removal of duplicates, 127 records were assessed for eligibility and 59 records subjected to full-text assessment. Six additional records which did not clearly report any associations between nutrients involved in one-carbon metabolism and DNA methylation were further excluded leaving 53 studies which are included in the qualitative analysis and 8 publications included in the meta-analysis. Study screening, eligibility and selection processes are shown in **Figure 1**.

### Characteristics of Studies Included

#### *Study design and background characteristics*

Summary and key findings of the studies included are provided in **Tables 2-6** <sup>13,20-24,31-34,39-81</sup>. Overall, data from 9561 adults with ages ranging from 18-85 years were included in this systematic review. Study participants were from 13 countries (USA, UK, Germany, Italy, the Netherlands, Sweden, Australia, Malaysia, Poland, China, Chile, Korea and Ireland). The majority (74.6%; n = 41) of studies involved both male and female participants. RCTs and intervention studies without randomization constitute 29.1% (n = 16) and 18.2% (n = 10) of studies reviewed respectively while 52.7% (n = 29) of the studies were observational (cross-section, case-control and cohort). While 1 publication reported both RCT and cross-sectional data <sup>23</sup>, another publication reported data for both an intervention and RCT <sup>40</sup>. Of the intervention studies included, 6 were depletion-repletion studies <sup>40,46,49,50,53,54</sup> and another 3 were supplementation studies without randomization <sup>47,51,52</sup>. Studies were conducted mainly in healthy individuals (34.6%, n = 19), or those with cancer (27.3%, n = 15), CVD (9.1%, n = 5), elderly subjects (9.1%, n = 5) and other diseases or conditions (20.0%, n = 11).

## 225 *One-carbon metabolism nutrients examined*

226           The main one-carbon metabolism nutrient supplemented or examined in most studies  
 227 (RCTs, intervention and observational) was folate (40.7%, n = 22), 18 studies (33.3%)  
 228 examined both folate/folic acid and vitamin B-12 status, 13 studies (24.1%) examined a  
 229 complex of B-vitamins and calcium and 1 study (1.9%) investigated methionine <sup>76</sup>. A large  
 230 proportion of studies measured biomarkers (75.0%, n = 39), 9 studies (17.3%) reported both  
 231 biomarker and dietary intake and 2 studies (3.8%) reported only dietary data. Of the 16  
 232 RCTs, 9 <sup>19,22,24,32–34,41,42,45</sup> supplemented with folic acid only (with doses ranging from  
 233 100µg/d to 1500µg/d). While 4 RCT studies intervened with a combination of folic acid and  
 234 vitamin B-12 <sup>21,23,32,44</sup>, another 4 RCT studies supplemented with folic acid and other B-  
 235 vitamins <sup>20,31,40,43</sup>. Duration of RCTs ranged from 10-156 weeks.

236           Furthermore, intervention studies using the depletion-repletion study design, fed  
 237 participants a folate-restricted diet (56µg-79.4µg/d) during the depletion stage and a folate  
 238 treatment diet (111µg-516µg/d) during the folate repletion stage. Observational studies  
 239 examined mainly circulating biomarker concentrations of folate and vitamin B-12 (51.7%, n  
 240 = 15), or circulating biomarker concentrations of several one-carbon metabolism nutrients  
 241 including folate, B-12, B-6, B-2, betaine, choline and methionine (34.5%, n = 10) and 4  
 242 studies (13.8%) examined only folate status.

## 243 *DNA methylation Analysis*

244           Studies focused on a range of genomic locations and DNA methylation was assessed  
 245 in a variety of tissues using different methods. While 67.3% of studies (n = 37) examined  
 246 global methylation, 20.0% (n = 11) measured gene-specific methylation, 1 study (1.8%)  
 247 examined genome-wide methylation <sup>21</sup> while 6 studies (10.9%) examined both global and  
 248 gene-specific methylation <sup>20,22,52,59,60,68</sup>. Methylation was examined mostly in blood (whole  
 249 blood, leukocytes, monocytes and peripheral blood cells; 74.6%, n = 41), colorectal tissue

(20.0%, n = 11) or both blood and colon tissues (5.5%, n = 3). DNA methylation analyses were carried out using 16 different techniques, mainly pyrosequencing (n = 14), LC-MS techniques (n = 12), methyl acceptance assay (n = 12) and 6 studies used more than one method.

## **Effect of Supplementation with Nutrients Involved in One-Carbon Metabolism on DNA Methylation**

The effect of supplementation with nutrients involved in one-carbon metabolism on DNA methylation was investigated in 16 studies using RCT study design<sup>19–24,31–34,40–45</sup> (**Table 2**). While the largest proportion of RCTs examined the effect of one-carbon metabolism nutrients on global methylation (68.8%, n = 11), 2 studies (12.5%) investigated gene-specific methylation<sup>44,45</sup>, another 2 studies (12.5%) examined both global and gene-specific methylation (12.5%) and 1 study (6.3%) examined genome-wide methylation. While 61.5% (n = 8) of RCTs examining global DNA methylation observed no significant changes in methylation in response to supplementation<sup>22–24,32,34,40,41,43</sup>, 38.5% (n = 5) observed significant increases in methylation<sup>19,20,31,33,42</sup>. Furthermore, RCTs investigating gene-specific methylation in colorectal adenoma patients and elderly subjects showed significant increases in colorectal tissue and blood DNA methylation at several loci including *ASPA*, *PDE4C*, *MGMT*, *MLH1*, *p14*, *p16* and *RASSF1A*<sup>20,44</sup> in response to nutrient supplementation; however no significant effects were observed for *ESR1*, *ITGA2B*, *MLH1* and *SFRP1* methylation<sup>20,22,45</sup>.

In the single RCT investigating the effects of supplementation with folic acid and vitamin B-12 on epigenome-wide methylation in adult leukocyte samples<sup>21</sup>, 6 significant differentially methylated regions (DMRs) between intervention and placebo groups were discovered. Intervention with folic acid and B-12 in this study increased DNA methylation for the majority of *HOX* genes while remaining stable or decreasing in the placebo group. In

addition to comparisons for DNA methylation changes between these two groups, the relationship between DNA methylation and serum folate was examined in a continuous manner, revealing that for 91% of the top 35 differentially methylated positions (DMPs), DNA methylation was positively correlated with levels of serum folate. Furthermore, 173 and 425 DMRs, were significantly associated (Benjamini-Hochberg adjusted p-value < 0.05) with serum folate and vitamin B-12 concentrations respectively in this study <sup>21</sup>.

Global methylation was examined in 90% (n = 9) of studies using an intervention study design without randomization with 1 study <sup>52</sup> investigating both global and gene-specific methylation (**Table 2**)<sup>40,46–51,53,54</sup>. Although 3 (60.0%) intervention studies conducted in both pre and postmenopausal women report decreased global methylation during folate restriction <sup>50,53,54</sup>, and 2 studies (40.0%) conducted in healthy premenopausal women observed no effect in response to depletion <sup>40,46</sup>. Conversely, in the intervention studies employing supplementation, 50.0% of these (n = 5) observed effects of supplementation on global methylation <sup>46–50,54</sup> and 50.0% (n = 5) did not observe any changes in methylation in both healthy populations or those with elevated homocysteine <sup>40,48,51,53</sup>. In the single intervention study investigating gene-specific methylation conducted in apparently healthy adults at increased risk of colorectal adenoma, there was no significant effect on methylation of 432 genes known to be abnormally methylated in human cancers <sup>52</sup>.

### **Association between Nutrients involved in One-Carbon Metabolism and DNA Methylation**

The majority of observational studies examined global methylation (56.7%, n=17), while 10 studies examined gene-specific methylation (33.3%) and three studies (10.0%) examined both (**Tables 3- 5**) <sup>13,23,55–81</sup>. Several observational studies indicate significant associations between nutrients and global methylation. While 9 studies (50.0%) direct associations <sup>58,66,70,71,76,78,81,82</sup>, 3 studies (16.7%) involving cancer patients or healthy

participants observed inverse correlations<sup>64,69,74</sup> and 6 studies (33.3%) did not observe any significant associations<sup>23,59,60,62,65,73</sup>. Two studies conducted in atherosclerosis patients and older participants measuring global methylation using 2 different surrogate markers of global methylation, observe positive associations between B-12 or B-6 status and Alu but not LINE-1 methylation<sup>63,67</sup>. A further 9 observational studies (75.0%) report significant associations between nutrients and methylation of specific gene loci with positive correlations observed for *VDR*, *p73*, *MTHFR*, *CACNA1G* and *RUNX3*<sup>55,72,75,77,79</sup> but negative correlations for *TNFA*, *MLH1*, *MGMT* and *ESR1* in cancer or obese patients<sup>13,56,68</sup>. Three studies (25.0%) did not observe significant correlations between nutrient status and *ec-SOD*, *p66Shc* and *TERT* methylation<sup>59,61,80</sup>.

### **Nutrients involved in One-Carbon Metabolism, Global Methylation and *MTHFR* C677T Genotype**

The *MTHFR* C677T polymorphism is a common polymorphism associated with reduced activity of the *MTHFR* enzyme and thereby affecting folate availability in one-carbon metabolism<sup>83</sup>. Sixteen studies examined the relations between nutrients involved in one-carbon metabolism (mainly folate), global methylation and the *MTHFR* C677T genotype. Low folate status was associated with lower methylation in *MTHFR* 677TT genotype participants compared to CC subjects<sup>58,66,82,84</sup>. Furthermore, decreases in methylation were observed in participants with the *MTHFR* 677TT genotype in response to folate supplementation in healthy young women<sup>41,46,54</sup>. On the contrary, 6 studies found no significant effect or association between folate status and DNA methylation in individuals stratified by the *MTHFR* C677T genotype<sup>24,40,44,48,60,62</sup>. Although stratification by *MTHFR* C677T genotype revealed 5 positions with differential methylation in response to B-vitamin supplementation, the power of the subgroup analysis was limited owing to low numbers and results should be interpreted with caution<sup>21</sup>.

## **Risk of Bias and Quality of Reporting Studies**

The quality of evidence presented in the included studies was rated as moderate with an average score of 65.2% on the quality assessment scales. Overall, RCT studies showed low risk of bias for random sequence generation, (93.8%), allocation concealment (75.0%), blinding of participants and personnel (75.0%), blinding of outcome assessment (37.5%), incomplete outcome data (43.8%) and selective reporting (37.5%) bias domains while all studies showed unclear risk of bias in other bias owing to lack of sufficient information to assess whether an important risk of bias exists. The majority of intervention studies showed an unclear risk of bias in all the domains owing to insufficient information provided to permit judgement (**Supplementary Table S3**). Observational studies showed high comparability and reporting of outcomes but were rated lower on the selection scale using the Newcastle-Ottawa scale (**Supplementary Table S4**). Although the quality of reporting studies was rated as good, several of the studies showed an unclear risk of bias highlighting the need for high quality studies with DNA methylation as the primary outcome.

## **Meta-Analysis on the Effect of Supplementation with Nutrients Involved in One-Carbon metabolism on Global DNA Methylation**

The meta-analysis examined the effect of supplementation with one-carbon metabolism nutrients on global DNA methylation. It included data pooled from 918 individuals across 8 RCT studies. Firstly, 9 publications were considered for inclusion in the meta-analysis. Of these, 1 study<sup>43</sup> was excluded through lack of numerical data and although attempts to contact the author was made, the data could not be obtained. A study reported post-supplementation data at 2 time points<sup>23</sup>, 1 study reported methylation data for both leukocytes and colon tissue<sup>19</sup> and another study<sup>32</sup> reported the effect of folic acid and a combination of folic acid and vitamin B-12 separately on global methylation. Although median values of methylation were reported in Kim *et al.*<sup>42</sup>, the corresponding dispersion



was not available. For the purposes of the meta-analysis, the SD value was extrapolated from Fenech *et al.*<sup>23</sup>, as similar methods were used for examining DNA methylation and methylation values were expressed in the same units.

Meta-analyses using the random effects model (**Figure 2**)<sup>19,20,23,24,32–34,42</sup> showed no significant overall effect of one-carbon metabolism nutrients on global DNA methylation ( $Z = 0.03$ ,  $P = 0.98$ ;  $I^2 = 64\%$ ,  $P = 0.002$ ). Pre-specified subgroup analyses of methylation in blood and colorectal tissue also indicated no effect of nutrient supplementation on global methylation in either blood ( $Z = 0.28$ ,  $P = 0.78$ ) or colon ( $Z = 0.60$ ,  $P = 0.55$ ). Substantial heterogeneity was observed in blood ( $I^2 = 71\%$ ,  $P = 0.002$ ,  $n = 7$ ) and non-significant moderate heterogeneity was observed for colorectal tissue ( $I^2 = 48\%$ ,  $P = 0.13$ ,  $n = 4$ ) in subgroup analyses.

Further pre-specified subgroup analysis focusing on the assay used to quantify global DNA methylation was carried out to attempt to explain the substantial heterogeneity among studies (**Figure 3**)<sup>20,23,24,32,34,39,42</sup>. Studies that assessed global DNA methylation by LC-MS techniques showed that B-vitamin supplementation significantly increased global DNA methylation ( $Z = 3.31$ ,  $P = 0.0009$ ). This finding was in contrast to the results of the individual studies, which did not find a significant effect of B-vitamin supplementation on DNA methylation. There was no detectable effect in studies using pyrosequencing ( $Z = 0.40$ ,  $P = 0.69$ ) and methyl acceptance assay ( $Z = 0.18$ ,  $P = 0.85$ ). No heterogeneity was observed for studies employing LC-MS techniques ( $I^2 = 0\%$ ,  $P = 0.60$ ,  $n = 3$ ) compared to pyrosequencing ( $I^2 = 76\%$ ,  $P = 0.04$ ,  $n = 2$ ) and methyl acceptance assay ( $I^2 = 64\%$ ,  $P = 0.03$ ,  $n = 5$ ). When analyses were focused on intervention with either folic acid or combination of B-vitamins (**Supplementary Figure S1**)<sup>20,23,24,32–34,39,42</sup>, subgroup analysis indicated no significant effect on DNA methylation owing to supplementation with folic acid only ( $Z = 0.52$ ,  $P = 0.60$ ) or folic acid in combination with B-12 and B-6 ( $Z = 0.52$ ,  $P = 0.61$ ).

Substantial heterogeneity was observed in the subgroup supplemented with folic acid only ( $I^2 = 71.0\%$ ,  $P = 0.002$ ,  $n = 7$ ) and B-vitamin combination ( $I^2 = 56\%$ ,  $P = 0.08$ ,  $n = 4$ ).

### **Publication Bias**

Publication bias assessment by visual inspection of the funnel plot did not indicate any substantial asymmetry and this was confirmed by a non-significant Egger's regression test ( $P = 0.152$ ).

### **Sensitivity Analysis**

A sensitivity analysis was performed by omitting one study at a time and assessing the pooled effect (Z) for the remaining studies. The pooled overall effect was consistent and within an acceptable range of 0.04 ( $P = 0.97$ ) to 0.57 ( $P = 0.57$ ). These findings indicate that the overall effect and heterogeneity are not significantly influenced by any particular study included in the meta-analysis.

### **Functional Analysis of Epigenome-Wide Methylation**

Further functional enrichment analysis was carried out on 173 and 425 DMRs which were shown to be significantly related to serum folate and vitamin B-12 status respectively (BH-adjusted p-value  $< 0.05$ ) in the single epigenome-wide methylation RCT<sup>21</sup>. The present analysis highlighted a subset of 12 DMRs (based on the exact genomic coordinates) which were significantly associated with both serum folate and vitamin B-12 status (**Figure 4a**). These are referred to as overlapping DMRs. The list of genes mapped to these DMRs are listed in (**Supplementary Table S5**). The overlapping DMRs were located in the first exon, gene body, TSS200, TSS1500, 3'UTR and 5'UTR (**Figure 4b**).

## DISCUSSION

This study encompasses qualitative and quantitative data to provide comprehensive evidence for the significant relationship between nutrients involved in one-carbon metabolism and DNA methylation across a range of health outcomes. The results from this systematic review indicate a significant role for specific nutrients in modulating both global and gene-specific methylation in a wide spectrum of diseases. Additionally, meta-analysis of a predefined subset of RCTs stratified by analytical method showed a significant increase in global methylation in response to B-vitamin supplementation for studies employing sensitive LC-MS techniques ( $n = 3$ ). This functional relationship between one-carbon metabolism nutrients and global methylation has not been previously estimated in meta-analysis of randomized trials.

While limited by the small number of RCTs in the meta-analysis of the LC-MS subgroup, a small but significant increase in global DNA methylation following supplementation with folic acid alone or in combination with vitamin B-12 was detected in comparison to studies using LINE-1 pyrosequencing or the methyl acceptance assay. LC-MS is an extremely sensitive quantitative measure of total cellular 5-methyl cytosine<sup>85,86</sup> with guidelines<sup>87,88</sup> published on standardization and validation of methods by both the Food and Drug Administration (FDA) and European Medicines Agency (EMA). It is perhaps not surprising, therefore, that no heterogeneity was observed in the LC-MS subgroup analysis, which facilitated detection of the effect of B-vitamin supplementation on DNA methylation. Pyrosequencing of LINE-1 is also considered to be a very sensitive method and a multicenter benchmarking study evaluating DNA methylation techniques demonstrated that pyrosequencing of repetitive elements gave rise to highly reproducible results<sup>89</sup>. However, in the current studies reviewed, heterogeneity may have arisen through the use of non-standardized protocols in various laboratories, resulting in analysis of varying regions of the

LINE-1 locus or assays using a varied number of CpG sites and thereby preventing detection of a significant effect. Although providing a reasonable estimate of global methylation, the methyl acceptance assay is a semi-quantitative assay, confounded by suboptimal enzyme activity and stability of SAM, resulting in large assay variability.

Furthermore, it is established that DNA methylation is highly tissue-specific and variability in tissues analyzed could mask potential associations between specific nutrients and DNA methylation<sup>90</sup>. Each of the studies in the LC-MS subgroup analyzed methylation in a single tissue type, i.e. blood, while LINE-1 pyrosequencing and methyl acceptance assay studies were conducted using a mixture of both blood and colorectal tissue further confounding the meta-analysis of these groups. LINE-1 pyrosequencing is often the preferred method over LC-MS for global methylation as the necessary expertise and equipment for LC-MS are not as widely available<sup>91</sup>. In order to enable a meaningful comparison of global methylation between studies using LINE-1 pyrosequencing, there is a need for researchers in the field to adopt a more standardized approach. Given the already proven reproducibility of the assay employed in laboratories of the BLUEPRINT consortium<sup>89</sup>, a reasonable recommendation would therefore be the widespread adoption of this method.

Assessment of DNA methylation in the studies reviewed here covered a wide range of genomic regions using 16 different techniques, introducing substantial variability and leading to confounding of outcome measurements and thereby posing significant challenges for comparability. It is therefore perhaps not surprising that the current meta-analysis did not detect an overall effect of supplementation with one-carbon metabolism nutrients on global methylation. In contrast to the current findings, a recent meta-analysis<sup>92</sup> reported increased global DNA methylation in response to folic acid in colorectal mucosa but not blood. This result was, however, largely driven by a single study<sup>93</sup> that was shown by the authors to display publication bias. In agreement with the current investigation, reanalysis of the data

excluding this study did not detect a significant effect of folic acid on global methylation<sup>92</sup>. The study by Cravo *et al.*<sup>93</sup> did not in fact meet the strict inclusion criteria for the current systematic review and meta-analysis. These discrepancies in findings from both studies are explained by the limited number of RCTs which met the inclusion criteria for meta-analysis. This therefore highlights the urgent need for further high quality, robustly designed RCT studies of B-vitamin supplementation to identify epigenetic modification in response to B-vitamin supplementation.

Folic acid was the main nutrient used for supplementation in the majority of the RCTs included in the meta-analysis. It was also the main B-vitamin investigated in one-third of intervention and observational studies. By solely focusing on folate, interventions may only partly modify the dietary factors related to one-carbon metabolism that influence DNA methylation as the availability of folate depends on other B-vitamins<sup>24,94</sup>. Furthermore, the interactions between nutrients involved in one-carbon metabolism when supplemented in combination is complex, currently not fully understood and could influence DNA methylation through different mechanisms<sup>94</sup>. For example, in an RCT of long-term supplementation with folic acid and vitamin B-12 in elderly subjects, methylation of 425 DMRs is significantly associated with serum vitamin B-12 concentrations, whereas only 173 DMRs are associated with serum folate status<sup>21</sup>. Novel functional analysis presented here highlights a subset of only 12 DMRs significantly related to both serum folate and vitamin B-12 status. The genes mapped to these DMRs will be a valuable resource for future studies investigating the combined effects of B-vitamin supplementation on DNA methylation and may provide future targets for epigenetic therapies. Similarly, the effect of one-carbon metabolism nutrients on DNA methylation is influenced by specific gene mutations in the one-carbon pathway and polymorphisms (including the *MTHFR* C677T polymorphism) that affect availability of methyl groups for methylation reactions in one-carbon metabolism<sup>83</sup>.

More comprehensive studies are required to examine the complex interaction between polymorphisms in one-carbon metabolism, B-vitamins and other related nutrients in relation to DNA methylation.

Strengths of this current study include the use of mixed qualitative and quantitative approaches to provide comprehensive evidence of the relations between nutrients involved in one-carbon metabolism and DNA methylation. Additionally, the strength of the meta-analysis was that only RCTs with the most robust design (i.e. RCTs with a parallel placebo or control group) were included. A potential limitation of the present study is that a meaningful quantitative pooling of data could only be performed for a small subset of RCTs owing to substantial heterogeneity in study aims, designs, population and health status, DNA methylation analysis techniques and tissues analyzed. Further, while most studies investigating gene-specific methylation and one-carbon metabolism nutrients followed a candidate gene approach, providing a valuable starting point for future investigations, there is the possibility that other loci, yet to be identified, which may also be influenced by one-carbon metabolism nutrients could have been overlooked.

## CONCLUSION

In conclusion, the present systematic review supports a functional relationship between specific nutrients involved in one-carbon metabolism and DNA methylation. Meta-analysis, of the limited evidence currently available from RCTs, shows that supplementation with folic acid alone or in combination with vitamin B-12 resulted in an increase in global DNA methylation. The results of this study provide a foundation for further work, and highlight the need for future studies investigating the role of B-vitamins in epigenetic modifications associated with disease.

**Acknowledgements:** We thank the authors of the original articles, in particular those who were approached for additional information. Also Joan Atkinson, Ulster University Library for help with the systematic search.

**Authors' Contributions were as follows:**

DLM, CFH and MW designed the research; SDA, SR, JD conducted the research, SDA performed the statistical analysis; SDA, DEK and S-JT performed the functional analysis; SDA, CFH, MW and DLM wrote the article; HM, JJS and CPW carried out critical revision for important intellectual content. DLM had primary responsibility for the final content. All authors read and approved the final version of the manuscript.

**Funding:** Northern Ireland Chest, Heart & Stroke Association (NICH206\_07) (DLM, MW), Vice Chancellor's Research Scholarship (SDA), Ulster University. The funders played no role in the conception, design, performance, and approval of the work.

**Declaration of interest:** None of the authors have relevant interests to declare.

**Supporting information:** The following supporting information is available through the online version of this article at the publisher's website:

**Figure S1:** Meta-analysis of the effect of supplementation with nutrients involved in one-carbon metabolism on global DNA methylation sub-grouped by one-carbon metabolism nutrients.

**Table S1:** PRISMA checklist

**Table S2:** Systematic search strategy

**Supplementary Table S3:** Risk of bias assessment to randomized controlled trials and intervention studies investigating the effect nutrients involved in one-carbon metabolism on DNA methylation.

**Supplementary Table S4:** Risk of bias assessment of observational studies association between nutrients involved in one-carbon metabolism and DNA methylation

- 523 **Table S5:** Functional analysis of overlapping differentially methylated regions (DMRs)  
 524 related to both serum folate and vitamin B12 levels in epigenome-wide methylation analysis  
 525 **Table S6:** Full list of genes investigated in gene-specific methylation studies

## REFERENCES

1. Dor Y, Cedar H. Principles of DNA methylation and their implications for biology and medicine. *Lancet*. 2018;392:777-786. doi:10.1016/S0140-6736(18)31268-6
2. Miao L, Yin R-X, Zhang Q-H, et al. Integrated DNA methylation and gene expression analysis in the pathogenesis of coronary artery disease. *Aging (Albany NY)*. 2019;11(5):1486-1500.
3. Lu AT, Quach A, Wilson JG, et al. DNA methylation GrimAge strongly predicts lifespan and healthspan. *Aging (Albany NY)*. 2019;11(2):303-326.
4. Horvath S, Raj K. DNA methylation-based biomarkers and the epigenetic clock theory of ageing. *Nat Rev Genet*. 2018;19:371-384. doi:10.1038/s41576-018-0004-3
5. Madrigano J, Baccarelli A, Mittleman MA, et al. Aging and epigenetics: Longitudinal changes in gene-specific DNA methylation. *Epigenetics*. 2012;7(1):63-70. doi:10.4161/epi.7.1.18749
6. Marioni RE, Suderman M, Chen BH, et al. Tracking the Epigenetic Clock Across the Human Life Course: A Meta-analysis of Longitudinal Cohort Data. *J Gerontol A Biol Sci Med Sci*. 2019;74(1):57-61. doi:10.1093/gerona/gly060
7. Mandaviya PR, Joehanes R, Brody J, et al. Association of dietary folate and vitamin B-12 intake with genome-wide DNA methylation in blood: a large-scale epigenome-wide



- association analysis in 5841 individuals B-vitamin intake and genome-wide DNA methylation. *Am J Clin Nutr.* 2019;3:1-14. doi:10.1093/ajcn/nqz031/5511461
8. Martin EM, Fry RC. Environmental Influences on the Epigenome: Exposure-Associated DNA Methylation in Human Populations. *Annu Rev Public Heal.* 2018;39:309-333. doi:10.1146/annurev-publhealth
  9. Hannon E, Knox O, Sugden K, et al. Characterizing genetic and environmental influences on variable DNA methylation using monozygotic and dizygotic twins. *PLoS Genet.* 2018;14(8):1-27. doi:10.1371/journal.pgen.1007544
  10. Altmann S, Murani E, Schwerin M, Metges CC, Wimmers K, Ponsuksili S. Somatic cytochrome c (CYCS) gene expression and promoter-specific DNA methylation in a porcine model of prenatal exposure to maternal dietary protein excess and restriction. *Br J Nutr.* 2012;107:791-799. doi:10.1017/S0007114511003667
  11. Dominguez-Salas P, Moore SE, Cole D, et al. DNA methylation potential: dietary intake and blood concentrations of one-carbon metabolites and cofactors in rural African women. *Am J Clin Nutr.* 2013;97:1217-1227. doi:10.3945/ajcn.112.048462
  12. Steegers-Theunissen RP, Obermann-Borst SA, Kremer D, et al. Periconceptional Maternal Folic Acid Use of 400 mg per Day Is Related to Increased Methylation of the IGF2 Gene in the Very Young Child. *PLoS One.* 2009;4(11):1-5. doi:10.1371/journal.pone.0007845
  13. Bollati V, Favero C, Albetti B, et al. Nutrients intake is associated with DNA methylation of candidate inflammatory genes in a population of obese subjects. *Nutrients.* 2014;6(10):4625-4639. doi:10.3390/nu6104625
  14. Malcomson FC, Mathers JC. Nutrition, epigenetics and health through life. *Nutr Bull.* 2017;42(3):254-265. doi:10.1111/nbu.12281
  15. Glier MB, Green TJ, Devlin AM. Methyl nutrients, DNA methylation, and

- cardiovascular disease. *Mol Nutr Food Res*. 2014;58(1):172-182.  
doi:10.1002/mnfr.201200636
16. Waterland RA, Kellermayer R, Laritsky E, et al. Season of Conception in Rural Gambia Affects DNA Methylation at Putative Human Metastable Epialleles. *PLoS Genet*. 2010;6(12):1-10. doi:10.1371/journal.pgen.1001252
  17. Tobi EW, Lumey LH, Talens RP, et al. DNA methylation differences after exposure to prenatal famine are common and timing- and sex-specific. *Hum Mol Genet*. 2009;18(21):4046-4053. doi:10.1093/hmg/ddp353
  18. Mason JB. Biomarkers of Nutrient Exposure and Status in One-Carbon (Methyl) Metabolism. *J Nutr*. 2003;133.
  19. Pufulete M. Effect of folic acid supplementation on genomic DNA methylation in patients with colorectal adenoma. *Gut*. 2005;54:648-653. doi:10.1136/gut.2004.054718
  20. Obeid R, Hübner U, Bodis M, Graeber S, Geisel J. Effect of adding B-vitamins to vitamin D and calcium supplementation on CpG methylation of epigenetic aging markers. *Nutr Metab Cardiovasc Dis*. 2018;28(4):411-417.  
doi:10.1016/j.numecd.2017.12.006
  21. Kok DEG, Dhonukshe-Rutten RA, Lute C, et al. The effects of long-term daily folic acid and vitamin B 12 supplementation on genome- wide DNA methylation in elderly subjects. *Clin Epigenetics*. 2015;7(121):1-14. doi:10.1186/s13148-015-0154-5
  22. Al-Ghnaniem Abbadi R, Emery P, Pufulete M. Short-term folate supplementation in physiological doses has no effect on ESR1 and MLH1 methylation in colonic mucosa of individuals with adenoma. *J Nutrigenet Nutrigenomics*. 2013;5(6):327-338.  
doi:10.1159/000345819
  23. Fenech M, Aitken C, Rinaldi J. Folate, vitamin B12, homocysteine status and DNA damage in young Australian adults. *Carcinogenesis*. 1998;19(7):1163-1171.

24. Jung AY, Smulders Y, Verhoef P, et al. No Effect of Folic Acid Supplementation on Global DNA Methylation in Men and Women with Moderately Elevated Homocysteine. *PLoS One*. 2011;6(9). doi:10.1371/journal.pone.0024976
25. Higgins JPT, Altman DG. *Chapter 8: Assessing Risk of Bias in Included Studies.*; 2017.
26. Wells GA, Shea B, O'Connell D, et al. The Newcastle-Ottawa Scale (NOS) for assessing the quality of nonrandomised studies in meta-analyses. The Ottawa Hospital Research Institute.  
  
[http://www.ohri.ca/programs/clinical\\_epidemiology/nos\\_manual.pdf](http://www.ohri.ca/programs/clinical_epidemiology/nos_manual.pdf). Published 2011.
27. Vandenberg JP, Von Elm E, Altman DG, et al. Strengthening the Reporting of Observational Studies in Epidemiology (STROBE): Explanation and Elaboration. *Plos Med*. 2007;4(10). doi:10.1371/journal.pmed
28. Moher D, Hopewell S, Schulz KF, et al. CONSORT 2010 Explanation and Elaboration: updated guidelines for reporting parallel group randomised trials. *BMJ Open*. 2010;340(c869). doi:10.1136/bmj.c869
29. Jarlais DC Des, Lyles C, Crepaz N. Improving the Reporting Quality of Nonrandomized Evaluations of Behavioral and Public Health Interventions: The TREND Statement. *Am J Public Health*. 2004;94(3):361-366.
30. Dullemeijer C, Souverein O. *Guidance Document for: Meta-Analyses on RCTs with Continuous Outcome Variables.*; 2011.
31. Pusceddu I, Herrmann M, Kirsch SH, et al. Prospective study of telomere length and LINE-1 methylation in peripheral blood cells: the role of B vitamins supplementation. *Eur J Nutr*. 2016;55:1863-1873. doi:10.1007/s00394-015-1003-1
32. Stopper H, Treutlein AT, Bahner U, et al. Reduction of the genomic damage level in haemodialysis patients by folic acid and vitamin B12 supplementation. *Nephrol Dial*

- Transplant.* 2008;23(10):3272-3279. doi:10.1093/ndt/gfn254
33. O'Reilly SL, McGlynn AP, McNulty H, et al. Folic Acid Supplementation in Postpolypectomy Patients in a Randomized Controlled Trial Increases Tissue Folate Concentrations and Reduces Aberrant DNA Biomarkers in Colonic Tissues Adjacent to the Former Polyp Site. *J Nutr.* 2016;146(5):933-939. doi:10.3945/jn.115.222547
  34. Figueiredo JC, Grau M V., Wallace K, et al. Global DNA hypomethylation (LINE-1) in the normal colon and lifestyle characteristics and dietary and genetic factors. *Cancer Epidemiol Biomarkers Prev.* 2009;18(4):1041-1049. doi:10.1158/1055-9965.EPI-08-0926
  35. Higgins J, Green S. *Cochrane Handbook for Systematic Reviews of Interventions Version 5.1.0.*; 2011.
  36. Dersimonian R, Laird N. Meta-Analysis in Clinical Trials. *Control Clin Trials.* 1986;7:177-188.
  37. Egger M, Smith GD, Schneider M, Minder C. Papers Bias in meta-analysis detected by a simple, graphical test. *BMJ.* 1997;315:629-631.
  38. Hulsen T, De Vlieg J, Alkema W. BioVenn-a web application for the comparison and visualization of biological lists using area-proportional Venn diagrams. *BMC Genomics.* 2008;9(488):1-6. doi:10.1186/1471-2164-9-488
  39. Pufulete M, Al-Ghnaniem R, Khushal A, et al. Effect of folic acid supplementation on genomic DNA methylation in patients with colorectal adenoma. *Gut.* 2005;5(4):648-653. doi:10.1136/gut.2004.054718
  40. Abratte CM, Wang W, Li R, Axume J, Moriarty DJ, Caudill MA. Choline status is not a reliable indicator of moderate changes in dietary choline consumption in premenopausal women. *J Nutr Biochem.* 2009;20(1):62-69. doi:10.1016/j.jnutbio.2007.12.002

41. Crider KS, Quinlivan EP, Berry RJ, et al. Genomic DNA Methylation Changes in Response to Folic Acid Supplementation in a Population-Based Intervention Study among Women of Reproductive Age. Xu G, ed. *PLoS One*. 2011;6(12):e28144. doi:10.1371/journal.pone.0028144
42. Kim YI, Baik HW, Fawaz K, et al. Effects of folate supplementation on two provisional molecular markers of colon cancer: A prospective, randomized trial. *Am J Gastroenterol*. 2001;96(1):184-195. doi:10.1016/S0002-9270(00)02267-X
43. Nanayakkara PWB, Kiefte-De Jong JC, Stehouwer CDA, et al. Association between global leukocyte DNA methylation, renal function, carotid intima-media thickness and plasma homocysteine in patients with stage 2-4 chronic kidney disease. *Nephrol Dial Transplant*. 2008;23(8):2586-2592. doi:10.1093/ndt/gfn040
44. Van den Donk M, Pellis L, Crott JW, et al. Folic acid and vitamin B-12 supplementation does not favorably influence uracil incorporation and promoter methylation in rectal mucosa DNA of subjects with previous colorectal adenomas. *J Nutr*. 2007;137(9):2114-2120.
45. Wallace K, Grau M V., Levine AJ, et al. Association between folate levels and CpG island hypermethylation in normal colorectal mucosa. *Cancer Prev Res*. 2010;3(12):1552-1564. doi:10.1158/1940-6207.CAPR-10-0047
46. Axume J, Smith SS, Pogribny IP, Moriarty DJ, Caudill MA. The methylenetetrahydrofolate reductase 677TT genotype and folate intake interact to lower global leukocyte DNA methylation in young Mexican American women. *Nutr Res*. 2007;27:13-17. doi:10.1016/j.nutres.2006.12.006
47. Ellingrod VL, Grove TB, Burghardt KJ, Taylor SF, Dalack G. The effect of folate supplementation and genotype on cardiovascular and epigenetic measures in schizophrenia subjects. *npj Schizophr*. 2015;1(1):15046. doi:10.1038/npjschz.2015.46

48. Hubner U, Geisel J, Kirsch SH, et al. Effect of 1 year B and D vitamin supplementation on LINE-1 repetitive element methylation in older subjects. *Clin Chem Lab Med*. 2013;51(3):649-655. doi:10.1515/cclm-2012-0624
49. Ingrosso D, Cimmino A, Perna AF, et al. Folate treatment and unbalanced methylation and changes of allelic expression induced by hyperhomocysteinaemia in patients with uraemia. *Lancet*. 2003;1693-1699. doi:10.1016/S0140-6736(03)13372-7
50. Jacob R a, Gretz DM, Taylor PC, et al. Moderate folate depletion increases plasma homocysteine and decreases lymphocyte DNA methylation in postmenopausal women. *J Nutr*. 1998;128(7):1204-1212.
51. Pizzolo F, Henk BJ, Choi SW, et al. Folic Acid Effects on S-Adenosylmethionine, S-Adenosylhomocysteine, and DNA Methylation in Patients with Intermediate Hyperhomocysteinemia. *J Am Coll Nutr*. 2011;30(1):11-18.  
doi:10.1080/07315724.2011.10719939
52. Protiva P, Mason JB, Liu Z, et al. Altered folate availability modifies the molecular environment of the human colorectum: Implications for colorectal carcinogenesis. *Cancer Prev Res*. 2011;4(4):530-543. doi:10.1158/1940-6207.CAPR-10-0143
53. Rampersaud GC, Kauwell GP, Hutson AD, Cerda JJ, Bailey LB. Genomic DNA methylation decreases in response to moderate folate depletion in elderly women. *Am J Clin Nutr*. 2000;72:998-1003.
54. Shelnutt KP, Kauwell GPA, Gregory JF, et al. Methylenetetrahydrofolate reductase 677C→T polymorphism affects DNA methylation in response to controlled folate intake in young women. *J Nutr Biochem*. 2004;15(9):554-560.  
doi:10.1016/j.jnutbio.2004.04.003
55. Beckett EL, Duesing K, Martin C, et al. Relationship between methylation status of vitamin D-related genes, vitamin D levels, and methyl-donor biochemistry. *J Nutr*

- Intermed Metab.* 2016;6:8-15. doi:10.1016/j.jnim.2016.04.010
56. Coppedè F, Migheli F, Lopomo A, et al. Gene promoter methylation in colorectal cancer and healthy adjacent mucosa specimens: Correlation with physiological and pathological characteristics, and with biomarkers of one-carbon metabolism. *Epigenetics.* 2014;9(4):621-633. doi:10.4161/epi.27956
  57. Friso S, Choi S-W, Girelli D, et al. A common mutation in the 5,10-methylenetetrahydrofolate reductase gene affects genomic DNA methylation through an interaction with folate status. *Proc Natl Acad Sci U S A.* 2002;99(8):5606-5611. doi:10.1073/pnas.062066299
  58. Friso S, Girelli D, Trabetti E, et al. The MTHFR 1298A>C polymorphism and genomic DNA methylation in human lymphocytes. *Cancer Epidemiol Biomarkers Prev.* 2005;14(4):938-943. doi:14/4/938 [pii]r10.1158/1055-9965.EPI-04-0601
  59. Geisel J, Schorr H, Bodis M, et al. The vegetarian lifestyle and DNA methylation. *Clin Chem Lab Med.* 2005;43(10):1164-1169. doi:10.1515/CCLM.2005.202
  60. Hanks J, Ayed I, Kukreja N, et al. The association between MTHFR 677C>T genotype and folate status and genomic and gene-specific DNA methylation in the colon of individuals without colorectal neoplasia. *Am J Clin Nutr.* 2013;98:1564-1574. doi:10.3945/ajcn.113.061432
  61. Hirsch S, Ronco AM, Guerrero-Bosagna C, et al. Methylation status in healthy subjects with normal and high serum folate concentration. *Nutrition.* 2008;24(11-12):1103-1109. doi:10.1016/j.nut.2008.05.018
  62. Kok RM, Smith DEC, Barto R, et al. Global DNA methylation measured by liquid chromatography-tandem mass spectrometry: analytical technique, reference values and determinants in healthy subjects. *Clin Chem Lab Med.* 2007;45(7):903-911. doi:10.1515/CCLM.2007.137

63. Perng W, Villamor E, Shroff MR, et al. Dietary intake, plasma homocysteine, and repetitive element DNA methylation in the Multi-Ethnic Study of Atherosclerosis (MESA). *Nutr Metab Cardiovasc Dis*. 2014;24:614-622.  
doi:10.1016/j.numecd.2013.11.011
64. Pufulete M, Al-Ghnaniem R, Rennie J, et al. Influence of folate status on genomic DNA methylation in colonic mucosa of subjects without colorectal adenoma or cancer. *Br J Cancer*. 2005;92:838-842. doi:10.1038/sj.bjc.6602439
65. Stenvinkel P, Karimi M, Johansson S, et al. Impact of inflammation on epigenetic DNA methylation - A novel risk factor for cardiovascular disease? *J Intern Med*. 2007;261(5):488-499. doi:10.1111/j.1365-2796.2007.01777.x
66. Stern LL, Mason JB, Selhub J, Choi SW. Genomic DNA hypomethylation, a characteristic of most cancers, is present in peripheral leukocytes of individuals who are homozygous for the C677T polymorphism in the Methylenetetrahydrofolate reductase gene. *Cancer Epidemiol Biomarkers Prev*. 2000;9(8):849-853.
67. Wernimont SM, Clark AG, Stover PJ, et al. Folate network genetic variation, plasma homocysteine, and global genomic methylation content: a genetic association study. *BMC Med Genet*. 2011;12(150):1-10. doi:10.1186/1471-2350-12-150
68. Al-Ghnaniem R, Peters J, Foresti R, Heaton N, Pufulete M. Methylation of estrogen receptor alpha and mutL homolog 1 in normal colonic mucosa: association with folate and vitamin B-12 status in subjects with and without colorectal neoplasia. *Am J Clin Nutr*. 2007;86(4):1064-1072. doi:86/4/1064 [pii]
69. Badiga S, Siddiqui NR, Macaluso M, Johanning GL, Piyathilake CJ. Homocysteinemia is Associated with a Lower Degree of PBMC LINE-1 Methylation and a Higher Risk of CIN 2C in the U.S. Post-Folic Acid Fortification Era. *Nutr Cancer*. 2016;68(3):446-455. doi:10.1080/01635581.2016.1152388



70. Bednarska-Makaruk M, Graban A, Sobczyńska-Malefora A, et al. Homocysteine metabolism and the associations of global DNA methylation with selected gene polymorphisms and nutritional factors in patients with dementia. *Exp Gerontol.* 2016;81:83-91. doi:10.1016/j.exger.2016.05.002
71. Friso S, Udali S, Guarini P, et al. Global DNA hypomethylation in peripheral blood mononuclear cells as a biomarker of cancer risk. *Cancer Epidemiol Biomarkers Prev.* 2013;22(3):348-355. doi:10.1158/1055-9965.EPI-12-0859
72. Kim JW, Park HMi, Choi Y, Chong SY, Oh D, Kim NK. Polymorphisms in genes involved in folate metabolism and plasma DNA methylation in colorectal cancer patients. *Oncol Rep.* 2011;25:167-172. doi:10.3892/or
73. Nan H, Giovannucci EL, Wu K, et al. Pre-Diagnostic Leukocyte Genomic DNA Methylation and the Risk of Colorectal Cancer in Women. *PLoS One.* 2013;8(4). doi:10.1371/journal.pone.0059455
74. Pufulete M, Al-Ghnaniem R, Leather AJM, et al. Folate status, genomic DNA hypomethylation, and risk of colorectal adenoma and cancer: A case control study. *Gastroenterology.* 2003;124(5):1240-1248. doi:10.1016/S0016-5085(03)00279-8
75. Tannorella P, Stoccoro A, Tognoni G, et al. Methylation analysis of multiple genes in blood DNA of Alzheimer's disease and healthy individuals. *Neurosci Lett.* 2015;600:143-147. doi:10.1016/j.neulet.2015.06.009
76. Tremolizzo L, Messina P, Conti E, et al. Whole-blood global DNA methylation is increased in amyotrophic lateral sclerosis independently of age of onset. *Amyotroph Lateral Scler Front Degener.* 2014;15(1-2):98-105. doi:10.3109/21678421.2013.851247
77. Van Guelpen B, Dahlin AM, Hultdin J, et al. One-carbon metabolism and CpG island methylator phenotype status in incident colorectal cancer: A nested case-referent

- study. *Cancer Causes Control*. 2010;21(4):557-566. doi:10.1007/s10552-009-9484-y
78. Wang TC, Song YS, Wang H, et al. Oxidative DNA damage and global DNA hypomethylation are related to folate deficiency in chromate manufacturing workers. *J Hazard Mater*. 2012;213-214:440-446. doi:10.1016/j.jhazmat.2012.02.024
  79. Wei LK, Sutherland H, Au A, et al. A Potential Epigenetic Marker Mediating Serum Folate and Vitamin B 12 Levels Contributes to the Risk of Ischemic Stroke. *Biomed Res Int*. 2015;2015:1-4. doi:10.1155/2015/167976
  80. Zhang D, Wen X, Zhang L, Cui W. DNA Methylation of Human Telomerase Reverse Transcriptase Associated With Leukocyte Telomere Length Shortening in Hyperhomocysteinemia-Type Hypertension in Humans and in a Rat Model. *Circ J*. 2014;78(8):1915-1923. doi:10.1253/circj.CJ-14-0233
  81. Bae S, Ulrich CM, Bailey LB, et al. Impact of folic acid fortification on global DNA methylation and one-carbon biomarkers in the Women's Health Initiative Observational Study cohort. *Epigenetics*. 2014;9(3):396-403.
  82. Friso S, Choi S-W, Girelli D, et al. A common mutation in the 5,10-methylenetetrahydrofolate reductase gene affects genomic DNA methylation through an interaction with folate status. *PNAS*. 2002;99(8).
  83. Fredriksen A, Meyer K, Ueland PM, Vollset SE, Grotmol T, Schneede J. Large-Scale Population-Based Metabolic Phenotyping of Thirteen Genetic Polymorphisms Related to One-Carbon Metabolism. *Hum Mutat*. 2007;28(9):856-865. doi:10.1002/humu
  84. Friso S, Udali S, Guarini P, et al. Global DNA Hypomethylation in Peripheral Blood Mononuclear Cells as a Biomarker of Cancer Risk. *Cancer Epidemiol Biomarkers Prev*. 2013;22(3):348-355. doi:10.1158/1055-9965.EPI-12-0859
  85. Quinlivan EP, Gregory J. DNA methylation determination by liquid chromatography-tandem mass spectrometry using novel biosynthetic [U-15N]deoxycytidine and [U-15

- N]methyldeoxycytidine internal standards. *Nucleic Acids Res.* 2008;36(18):1-7.  
doi:10.1093/nar/gkn534
86. Liu J, Hesson LB, Ward RL. Liquid Chromatography Tandem Mass Spectrometry for the Measurement of Global DNA Methylation and Hydroxymethylation. *J Proteomics Bioinform.* 2013;2. doi:10.4172/jpb.S2-005
  87. Food and Drug Administration. *Bioanalytical Method Validation Guidance for Industry.*; 2018.
  88. European Medicines Agency. *Guideline on Bioanalytical Method Validation.*; 2011.
  89. Bock C, Halbritter F, Carmona FJ, et al. Quantitative comparison of DNA methylation assays for biomarker development and clinical applications. *Nat Biotechnol.* 2016;34(7):726-740. doi:10.1038/nbt.3605
  90. Houseman EA. DNA Methylation and Cell-Type Distribution. In: *Computational and Statistical Epigenomics*. Vol 7. ; 2015:35-50. doi:10.1007/978-94-017-9927-0
  91. Kurdyukov S, Bullock M, Ehrlich M. DNA Methylation Analysis: Choosing the Right Method. *Biology (Basel)*. 2016;5(3). doi:10.3390/biology5010003
  92. ElGendy K, Malcomson FC, Lara JG, Bradburn DM, Mathers JC. Effects of dietary interventions on DNA methylation in adult humans: systematic review and meta-analysis. *Br J Nutr.* 2018;120(9):961-976. doi:10.1017/S000711451800243X
  93. Cravo M, Fidalgo P, Pereira A. DNA methylation as an intermediate biomarker in colorectal cancer: modulation by folic acid supplementation. *Eur J Cancer Prev.* 1994;3:473-479.
  94. Caudill MA. Folate bioavailability: implications for establishing dietary recommendations and optimizing status. *Am J Clin Nutr.* 2010;91.  
doi:10.3945/ajcn.2010.28674E

## FIGURE LEGENDS

**FIGURE 1.** Flow diagram of study selection for systematic review and meta-analysis. <sup>1</sup>One publication reported data from both a cross-sectional study and RCT, another publication reported both RCT and intervention data.

**FIGURE 2.** Random-effects meta-analysis of the effect of supplementation with nutrients involved in one-carbon metabolism on global DNA methylation sub-grouped by tissues analyzed. The horizontal lines running through each square represent the 95% CI for each study. The diamonds indicate pooled effect and 95% CI for each subgroup and the overall effect (Z).  $\chi^2$ , chi-squared test assesses whether observed differences in results are compatible with chance alone;  $I^2$ , heterogeneity index (0–100%).

**FIGURE 3.** Random-effects meta-analysis of the effect of supplementation with one-carbon metabolism nutrients on global DNA methylation sub-grouped by methylation techniques. The horizontal lines running through each square represent the 95% CI for each study. The diamonds indicate pooled effect and 95% CI for each subgroup and the overall effect (Z).  $\chi^2$ , chi-squared test assesses whether observed differences in results are compatible with chance alone;  $I^2$ , heterogeneity index (0–100%).

**FIGURE 4.** Functional analysis of overlapping DMRs. a) A total of 12 DMRs were significantly associated with both folate and vitamin B-12 status in epigenome-wide studies. These DMRs are referred to as overlapping DMRs. b) Genomic locations of overlapping DMRs. TSS200, 200 base pairs around the transcription start site; TSS1500, 1500 base pairs around the transcription start site; 3' UTR, 3' untranslated region; 5' UTR, 5' untranslated region.

## TABLE LEGENDS

**TABLE 2.** Randomized controlled trials investigating the effect of one-carbon metabolism related nutrient supplementation on DNA methylation (n 16). <sup>a</sup>Full name of genes provided in Supplementary Table S6. **Abbreviations:** GC-MS, gas chromatography-mass spectrometry, LC-MS, liquid chromatography-tandem mass spectrometry

**TABLE 3.** Intervention studies investigating the effect of nutrients involved in one-carbon metabolism on DNA methylation (n 10). **Abbreviations:** LC/ESI-MS, liquid chromatography electrospray ionization mass spectrometry; LC-MS, liquid chromatography-tandem mass spectrometry, LUMA, luminometric assay; PBMC, peripheral blood mononuclear cells; MTHF, methyltetrahydrofolate.

**TABLE 4.** Cross-sectional studies investigating the association between nutrients involved in one-carbon metabolism and DNA methylation (n 15). <sup>a</sup>Full name of genes provided in Supplementary Table S6. **Abbreviations:** LC-MS, liquid chromatography-tandem mass spectrometry; LUMA, Luminometric methylation assay, PBMC - peripheral blood mononuclear cells.

**TABLE 5.** Case-control studies investigating the association between nutrients involved in one-carbon metabolism and DNA methylation (n 13). <sup>a</sup>Full name of genes provided in Supplementary Table S6. **Abbreviations:** LC-MS, liquid chromatography-tandem mass spectrometry; MS-HRM, methylation sensitive-high resolution melting analysis; PBMC, peripheral blood mononuclear cells.

**TABLE 6.** Cohort study investigating the association between nutrients involved in one-carbon metabolism and DNA methylation (n 1). **Abbreviations:** LC-MS, liquid chromatography-tandem mass spectrometry; RBC, red blood cell

**TABLE 1**

PICOS criteria for inclusion and exclusion of studies

<b>Parameter</b>	<b>Criteria</b>
Participants	Adults aged 18 years and older
Intervention	Supplementation with B-vitamins or dietary intake of B-vitamins
Comparison	Supplementation with nutrients involved in one-carbon metabolism compared to placebo or control group in the case of randomized controlled trials, pre/post same group comparison for intervention studies
Outcomes	Outcomes of interest were changes in DNA methylation (global, gene-specific, genome-wide) in response to supplementation with folic acid and related B-vitamins and association between dietary intake of one-carbon metabolism nutrients and DNA methylation (global, gene-specific, genome-wide).
Study design	Randomized and non-randomized intervention studies. Observational studies including cross-sectional, case-control and cohort studies

**TABLE 2**

Randomized controlled trials investigating the effect of one-carbon metabolism related nutrient supplementation on DNA methylation (n 16)

Study	Population	Country	n	Intervention		Duration	DNA methylation analysis			Main findings related to DNA methylation
				FA μg/d	Other μg/d or IU	Wk	Region	Technique	Tissue	
Global DNA methylation										
Abratte 2009 <sup>40</sup>	Heathy	USA	45	235.3		12	Global	Cytosine extension assay	Mononuclear cells	No effect
Abratte 2009 <sup>40</sup>	Heathy	USA	45	470.6	344000 choline 122000 betaine	12	Global	Cytosine extension assay	Mononuclear cells	No effect
Abratte 2009 <sup>40</sup>	Heathy	USA	45	235.3	412000 choline 267000 betaine	12	Global	Cytosine extension assay	Mononuclear cells	No effect
Abratte 2009 <sup>40</sup>	Heathy	USA	45	470.6	486000 choline 349000 betaine	12	Global	Cytosine extension assay	Mononuclear cells	No effect
Crider 2011 <sup>41</sup>	Healthy	China	135	100	-	24	Global	LC-MS	Blood	No effect
Crider 2011 <sup>41</sup>	Healthy	China	135	400	-	24	Global	LC-MS	Blood	No effect
Crider 2011 <sup>41</sup>	Healthy	China	135	4000	-	24	Global	LC-MS	Blood	No effect
Fenech 1998 <sup>23</sup>	Healthy	Australia	63	700	7 B-12	12	Global	Methyl acceptance assay	Lymphocyte	No effect
Fenech 1998 <sup>23</sup>	Healthy	Australia	63	2000	20 B-12	12	Global	Methyl acceptance assay	Lymphocyte	No effect
Figueiredo 2009 <sup>34</sup>	Colorectal adenoma	USA	388	1000	-	156	Global	Pyrosequencing	Colon	No effect
Jung 2011 <sup>24</sup>	Hyper-homocysteine	The Netherlands	216	800	-	156	Global	LC-MS	Leukocyte	No effect

Kim 2001 <sup>42</sup>	Adenoma	USA	20	5000	-	52	Global	Methyl acceptance assay	Colon	↑Methylation (p = 0.02)
Nanayakkara 2008 <sup>43</sup>	Chronic kidney disease	The Netherlands	78	5000	1000 B-6 1000 B-12	52	Global	LC-MS	Leukocyte	No effect
O'Reilly 2016 <sup>33</sup>	Adenoma	Ireland	20	600	-	34	Global	Modified alkaline comet assay	Colon	↑Methylation (p < 0.001)
Pufulete 2005 <sup>19</sup>	Colorectal adenoma	UK	33	400	-	10	Global	Methyl acceptance assay	Leukocyte Colon	↑Methylation in colonic mucosa (p = 0.09) leukocytes (p = 0.05)
Pusceddu 2016 <sup>31</sup>	Elderly subjects	Germany	60	500	500 B-12 50000 B-6 1200 vit D 456000 Ca	12	Global	Pyrosequencing	Whole blood	↑Methylation
Stopper 2008 <sup>32</sup>	Hemodialysis	Germany	27	6428.6	-	20	Global	LC-MS/MS	Whole blood	No effect
Stopper 2008 <sup>32</sup>	Hemodialysis	Germany	27	6428.6	142.9 B-12	20	Global	LC-MS/MS	Whole blood	No effect
<b><u>Gene-Specific Methylation</u></b>										
Van den Donk 2007 <sup>44</sup>	Colorectal adenoma	The Netherlands	81	4600	1100 B-12	24	<sup>a</sup> <i>MGMT, MLH1, p14, p16, APC, RASSF1A</i>	GC-MS MSP	Colorectal	↑Methylation (OR = 1.67, p = 0.08)
Wallace 2010 <sup>45</sup>	Colorectal adenoma	USA & Canada	388	1000	-	156	<i>ESR1, SFRP1</i>	Pyrosequencing	Colorectal	No effect
<b><u>Both Global and Gene-specific Methylation</u></b>										
Al-Ghnaniem Abbadi 2013 <sup>22</sup>	Colorectal adenoma	UK	29	400	-	10	Global <sup>a</sup> <i>ESR1, MLH1</i>	Methyl acceptance assay	Colon	No effect global, <i>ESR1, MLH1</i>



Pyrosequencing										
Obeid 2018 <sup>20</sup>	Elderly subjects	Germany	63	500	500 B-12, 50000 B-6 1200 vit D 456000 Ca	52	Global <sup>a</sup> <i>ASPA</i> , <i>ITGA2B</i> , <i>PDE4C</i>	Pyrosequencing	Whole blood	↑LINE-1 methylation ↑ <i>ASPA</i> methylation (p = 0.046) ↑ <i>PDE4C</i> methylation (p = 0.062) No effect <i>ITGA2B</i>
<b><u>Genome-wide DNA methylation</u></b>										
Kok 2015 <sup>21</sup>	Elderly subjects	The Netherlands	87	400	500 B-12	104	Genome-wide	Illumina 450k array	Buffly Coat	Differential methylation at 162 positions upon FA/vB-12 supplementation (1 DMP, cg19380919 sig) in intervention compared to placebo 6 DMRs differed between intervention and placebo groups Serum folate and vitamin B-12 significantly related to DNA methylation of 173 and 425 regions respectively.

<sup>a</sup>Full name of genes provided in **Supplementary Table S6**

Abbreviations: GC-MS, gas chromatography-mass spectrometry, LC-MS, liquid chromatography-tandem mass spectrometry

**TABLE 3**

Intervention studies investigating the effect of nutrients involved in one-carbon metabolism on DNA methylation (n 10)

Study	Population	Country	n	Intervention		Duration	DNA methylation analysis			Main findings related to DNA methylation
				FA μg/d	Other μg/d		Region	Technique	Tissue	
Global DNA Methylation										
Abratte 2009 <sup>40</sup>	Healthy	USA	45	78.24	344000 choline 122000 betaine	2	Global	Cytosine extension assay	PBMC	No effect
Abratte 2009 <sup>40</sup>	Healthy	USA	45	235.3	344000 choline 122000 betaine	12	Global	Cytosine extension assay	PBMC	No effect
Abratte 2009 <sup>40</sup>	Healthy	USA	45	470.6	344000 choline 122000 betaine	12	Global	Cytosine extension assay	PBMC	No effect
Abratte 2009 <sup>40</sup>	Healthy	USA	45	235.3	412000 choline 267000 betaine	12	Global	Cytosine extension assay	PBMC	No effect
Abratte 2009 <sup>40</sup>	Healthy	USA	45	470.6	486000 choline 349000 betaine	12	Global	Cytosine extension assay	PBMC	No effect
Axume 2007 <sup>46</sup>	Healthy	USA	43	79.4	-	12	Global	Cytosine extension assay	PBMC	No effect
Axume 2007 <sup>46</sup>	Healthy	USA	43	235.3	-	7	Global	Cytosine extension assay	PBMC	↓Methylation <i>MTHFR</i> 677TT (p < 0.05)
Axume 2007 <sup>46</sup>	Healthy	USA	43	470.6	-	7	Global	Cytosine extension assay	PBMC	↓Methylation <i>MTHFR</i> 677TT (p < 0.05)
Ellingrod 2015 <sup>47</sup>	Schizophrenia	USA	35	5000	-	12	Global	LUMA	Whole blood	↑Methylation (p < 0.0001)
Hubner 2013 <sup>48</sup>	Healthy	Germany	34	500	500 B-12 50000 B-6 1200IU vit D, 456000 Ca	52	Global	Pyrosequencing	Whole blood	No effect at 3 sites ↑Methylation at CpG site 317 (p = 0.044)
Ingrosso	Uremia/	Italy	14		15000 MTHF	8	Global	Cytosine	PBMC	↓Methylation

2003 <sup>49</sup>	Hyper-homocysteine							extension assay		
Jacob 1998 <sup>50</sup>	Healthy	USA	10	56	-	5	Global	Methyl acceptance assay	Lymphocyte	↓Methylation
Jacob 1998 <sup>50</sup>	Healthy	USA	10	111	-	4	Global	Methyl acceptance assay	Lymphocyte	↑Methylation
Jacob 1998 <sup>50</sup>	Healthy	USA	10	286	-	3	Global	Methyl acceptance assay	Lymphocyte	↑Methylation
Jacob 1998 <sup>50</sup>	Healthy	USA	10	516	-	3	Global	Methyl acceptance assay	Lymphocyte	↑Methylation
Pizzolo 2011 <sup>51</sup>	Moderate hyper-homocysteine	Italy	7	5000	-	8	Global	LC/ESI-MS	PBMC	No effect
Rampersaud 2002 <sup>53</sup>	Healthy	USA	33	118	-	7	Global	Methyl acceptance assay	Leukocyte	↓Methylation (p = 0.0025)
Rampersaud 2002 <sup>53</sup>	Healthy	USA	33	200	-	7	Global	Methyl acceptance assay	Leukocyte	No effect
Rampersaud 2002 <sup>53</sup>	Healthy	USA	33	415	-	7	Global	Methyl acceptance assay	Leukocyte	No effect
Shelnutt 2004 <sup>54</sup>	Healthy	USA	41	67.6	-	7	Global	Methyl acceptance assay	Leukocyte	↓Methylation (p = 0.08)
Shelnutt 2004 <sup>54</sup>	Healthy	USA	41	235	-	7	Global	LC/ESI-MS Methyl acceptance assay	Leukocyte	↑Methylation <i>MTHFR</i> 677TT (p < 0.05)
<b><u>Both Global and Gene-Specific</u></b>										
Protiva 2011 <sup>52</sup>	Healthy	USA	20	1000	-	8	Global Gene-specific	LC-MS Universal bead array	Colon	No effect

Abbreviations: LC/ESI-MS, liquid chromatography electrospray ionization mass spectrometry; LC-MS, liquid chromatography-tandem mass spectrometry, LUMA, luminometric assay; PBMC, peripheral blood mononuclear cells; MTHF, methyltetrahydrofolate

**TABLE 4**

Cross-sectional studies investigating the association between nutrients involved in one-carbon metabolism and DNA methylation (n 15)

Study	Population	Country	n	Nutrient Status		DNA methylation analysis			Main findings related to DNA methylation
				Biomarker	Dietary	Region	Technique	Tissue	
Global DNA Methylation									
Fenech 1998 <sup>23</sup>	Healthy	Australia	106	Folate, B-12	-	Global	Methyl acceptance assay	Lymphocyte	No correlation
Friso 2002 <sup>82</sup>	Valvular heart disease/ Healthy	Italy	292	Folate, B-12, B-6	-	Global	LC-MS	PBMC	Positive correlation with folate (p < 0.01), No correlation with B-12 status
Friso 2005 <sup>58</sup>	Healthy	Italy	198	Folate, B-12, B-6	-	Global	LC-MS	Lymphocyte	Positive correlation in <i>MTHFR</i> 1298 AA /677TT genotypes compared to the wild-type (p = 0.001)
Kok 2007 <sup>62</sup>	Healthy	The Netherlands	109	Folate, B-12, B-6, B2	-	Global	LC-MS	Blood	No correlation
Perng 2014 <sup>63</sup>	MESA study	USA	987	-	Folate, B-12, B-6, methionine	Global	Pyrosequencing	Leukocyte	Positive correlation with Alu No correlation with LINE-1
Pufulete 2005 <sup>64</sup>	Healthy	UK	68	Folate, B-12	Folate	Global	Methyl acceptance assay	Colon	Negative correlation serum folate (r = -0.311, p = 0.01), RBC folate (r = -0.356, p = 0.003), vitamin B-12 (r = -0.218, p = 0.08)
Stenvinkel 2007 <sup>65</sup>	Chronic kidney disease	Sweden	155	Folate, B-12	-	Global	LUMA	Leukocyte	No correlation

Stern 2000 <sup>66</sup>	Healthy	USA	19	Folate	-	Global	Methyl acceptance assay	Leukocyte	Positive correlation in <i>MTHFR</i> 677TT genotype (r= 0.738; p = 0.02)
Wernimont 2011 <sup>67</sup>	Normative ageing study	USA	621	Folate, B-12, B-6	-	Global	Pyrosequencing	Buffy coat	Correlation (p ≤ 0.05)
<b><u>Gene-specific DNA Methylation</u></b>									
Beckett 2016 <sup>55</sup>	Retirement health & lifestyle study	Australia	80	Folate, B-12	-	<sup>a</sup> <i>CY2R1, VDR, CYP27B1, CY24A1</i>	Epitect II methylation enzyme	Peripheral blood cells	Positive correlation ( <i>VDR</i> )
Bollati 2014 <sup>13</sup>	Obese/overweight	Italy	165		Folate, B-12	<sup>a</sup> <i>CD14, Et-1, iNOS, HERV-w, TNFα</i>	Pyrosequencing	Buffy coat	Negative correlation <i>TNFα</i> (β =-0.339, p = 0.012)
Coppede 2014 <sup>56</sup>	Colorectal cancer	Italy	107	Folate, B-12	-	<sup>a</sup> <i>APC, MGMT, MLH1, RASSF1A, CDKN2A, p16</i>	Methylation sensitive-high resolution melting	Tumor tissue	Negative correlation <i>MLH1</i> (p = 0.05)
Hirsch 2008 <sup>61</sup>	Healthy	Chile	111	Folate, B-12	-	<sup>a</sup> <i>ec-SOD</i>	Bisulfite sequencing	Lymphocyte	No correlation
<b><u>Both Global and Gene-specific DNA Methylation</u></b>									
Geisel 2005 <sup>59</sup>	Healthy	Germany	71	Folate, B-12, B-6	-	Global <sup>a</sup> <i>p66SHc</i>	Pyrosequencing	Whole blood	No correlation
Hanks 2013 <sup>60</sup>	Healthy	UK	336	Folate, B-12	Folate	Global <sup>a</sup> <i>ESR1, MYOD1, IGF2, N33, APC, MLH1, MGMT</i>	Pyrosequencing	Colon	No correlation global Negative correlation <i>MGMT</i> (p = 0.001)

<sup>a</sup>Full name of genes provided in **Supplementary Table S6**

Abbreviations: LC-MS, liquid chromatography-tandem mass spectrometry; LUMA, Luminometric methylation assay, PBMC - peripheral blood mononuclear cells

**TABLE 5**

Case-control studies investigating the association between nutrients involved in one-carbon metabolism and DNA methylation (n 13)

Study	Population	Country	n	Nutrient status			DNA methylation		Main findings related to DNA methylation
				Biomarker	Dietary	Region	Technique	Tissue	
Global DNA Methylation									
Badiga 2016 <sup>69</sup>	Cervical intraepithelial neoplasia	USA	132 case 325 control	Folate, B-12	-	Global	Pyrosequencing	PBMC	Negative correlation
Bednarska-Makaruk 2016 <sup>70</sup>	Dementia	Poland	102 case 45 control	Folate, B-12, 5-MTHF	-	Global	Imprint methylated kit	Leukocyte	Positive correlation (p = 0.013)
Friso 2013 <sup>71</sup>	Cancer	Italy	68 cancer 68 control	Folate	-	Global	LC-MS	PBMC	Positive correlation in <i>MTHFR</i> 677TT genotype
Nan 2013 <sup>73</sup>	Colorectal cancer	USA	358 CRC 661 control	Folate, B-12, B-6	Folate	Global	LC-MS	Leukocyte	No correlation
Pufulete 2003 <sup>74</sup>	Colorectal adenoma/cancer	UK	63 adenoma 76 control	Folate, B-12	Folate	Global	Methyl acceptance assay	Leukocyte/colon	Negative correlation Serum folate (r = -0.243, p = 0.009), RBC folate (r = -0.282, p = 0.002), folate status score (r = -0.295, p=0.001) in colon tissue No correlation in leukocytes
Tremolizzo 2013 <sup>76</sup>	ALS	Italy	96 ALS 87 control	Methionine	-	Global	Methyl acceptance assay	Whole blood	Positive correlation (r = 0.216, p = 0.043)
Wang 2012 <sup>78</sup>	Chromate exposure	China	115 case 60 control	Folate	-	Global	ELISA kit	Whole blood	Positive correlation (r = 0.163, p = 0.032)
Gene-Specific DNA Methylation									
Kim 2011 <sup>72</sup>	Colorectal cancer	Korea	67 CRC 53 control	Folate	-	<i>a</i> p16, p73, MLH1	Methylation-specific PCR	White blood cells	Positive correlation <i>p</i> 73

Tannorella 2015 <sup>75</sup>	Alzheimer's disease	Italy	120AD 115control	Folate, B- 12	-	<sup>a</sup> <i>PSEN1</i> , <i>BACE1</i> , <i>MTHFR</i> , <i>DNMT1</i> , <i>DNMT3A</i> , <i>DNMT3B</i> , <i>MTHFR</i>	Methylation MS-HRM	Peripheral blood	Positive correlation <i>MTHFR</i> methylation (r = 0.21; p = 0.002)
Van Guelpen 2009 <sup>77</sup>	Colorectal adenocarcino ma	Sweden		Folate, B- 12	-	<sup>a</sup> <i>CDKN2A</i> , <i>MLH1</i> , <i>IGF2</i> , <i>CACNA1G</i> , <i>NEUROG1</i> , <i>RUNX3</i> , <i>SOCS</i> , <i>CRABP1</i>	MethylLight real time PCR	Colon	Positive correlation vitamin B-12 <i>CACNA1G</i> (p = 0.047), folate <i>RUNX3</i> (p = 0.038)
Wei 2015 <sup>79</sup>	Ischemic stroke	Malaysia	297case 110 control	Folate, B- 12	-	<sup>a</sup> <i>MTHFR</i>	Pyrosequencing	Whole blood	Positive correlation Serum folate (r = 0.106, p = 0.032), vitamin B-12 (r = 0.114, p = 0.022)
Zhang 2014 <sup>80</sup>	Essential hypertension	China	258 EH 137 control	Folate, B- 12	-	<sup>a</sup> <i>TERT</i>	Methylation- specific PCR	Leukocyte	No correlation
<b><u>Both Global and Gene-Specific DNA Methylation</u></b>									
Al-Ghnaniem 2007 <sup>68</sup>	Colorectal neoplasia	UK	156 case 76 control	Folate, B- 12	-	Global <sup>a</sup> <i>ESR1</i> , <i>MLH1</i>	Methyl acceptance assay Pyrosequencing	Colon	Inverse association between serum folate/vitamin B12 and global methylation in adenoma patients. Negative correlation <i>ESR1</i> (r = 0.239, p = 0.003)

<sup>a</sup>Full name of genes provided in **Supplementary Table S6**

Abbreviations: LC-MS, liquid chromatography-tandem mass spectrometry; MS-HRM, methylation sensitive-high resolution melting analysis; PBMC, peripheral blood mononuclear cells

**TABLE 6**

Cohort study investigating the association between nutrients involved in one-carbon metabolism and DNA methylation (n 1)

Study	Population	Country	n	Nutrient Status		DNA methylation			Main findings related to DNA methylation
				Biomarker	Dietary	Region	Technique	Tissue	
Global DNA Methylation									
Bae 2014 <sup>81</sup>	Postmenopausal women	USA	408	Folate, B-12, B-6, choline, betaine	Folate, B-12, B-6, B2	Global	LC-MS	Leukocyte	Positive correlation Plasma folate (r = 0.20, p = 0.04) , RBC folate (r = 0.24, p = 0.01), vitamin B-12 (r = 0.18, p = 0.06)

Abbreviations: LC-MS, liquid chromatography-tandem mass spectrometry; RBC, red blood cell